

Geometric characteristics of conics in Bézier form

A. Cantón^a, L. Fernández-Jambrina^a, E. Rosado María^b

*^aDepartamento de Ciencias Aplicadas a la Ingeniería Naval
E.T.S.I. Navales
Arco de la Victoria s/n
E-28040-Madrid, Spain*

*^bDepartamento de Matemática Aplicada a la Edificación, al Medio Ambiente y al
Urbanismo
E.T.S. Arquitectura
Juan de Herrera 4
E-28040-Madrid, Spain*

Abstract

In this paper we address the calculation of geometric characteristics of conic sections (axes, asymptotes, centres, eccentricity, foci...) given in Bézier form in terms of their control polygons and weights, making use of real and complex projective and affine geometry and avoiding the use of coordinates.

Keywords: Conic sections, Bézier curves, eccentricity, asymptotes, axes, foci.

1. Introduction

Arcs of conic sections are easily represented as quadratic rational curves and this is the way they have been implemented in Computer Aided Geometric Design in exact form. Since they appear in almost every branch of industry, conic sections by themselves justify the inclusion of weights and denominators to extend polynomial curves in Bézier representation to rational ones.

Deriving closed formulas for geometric characteristics of conics given in Bézier form, that is, in terms of their control polygon and weights, was first accomplished in some cases in Lee (1987) by use of metric or Euclidean geometry. They may be found also in Piegl and Tiller (1997).

In Albrecht (2001) a different approach is followed, grounded in the calculation of the foci as intersections of the tangents to the conic from the

circular or absolute points. The main advantage of this approach is that the results are common for ellipses, hyperbolas and parabolas.

In Goldman and Wang (2004) closed formulas are obtained for computing geometric characteristics of conics from the invariants of rational quadratic parameterizations under rational linear reparameterizations instead.

On the other hand, in Sánchez-Reyes (2004) recipes are produced for constructing Bézier representations of conics when some of their geometric characteristics are known. In Sánchez-Reyes (2011) complex arithmetic is used to calculate some geometric characteristics of conic sections.

Most recently, in Xu et al. (2010) the eccentricity of conics in Bézier form is formulated and this result is used for deriving the range and extreme values of this parameter.

As a different approach we resort to projective and affine geometry to calculate geometric characteristics of conic sections in Bézier form. Some concepts such as center, foci and asymptotes can be defined at this level without introducing additional structure (scalar product, angles, lengths), though other geometric characteristics, such as axes, semi-axis lengths and eccentricity require calculation of angles and lengths and require therefore resorting to Euclidean geometry. The novel point in our approach is the possibility of writing the implicit equation of the conic in a coordinate-free expression that just uses the vertices and sides of the control polygon and the weights. This enables us to write geometric characteristics in closed form in a coordinate-free fashion.

In Section 2 we provide a quick review of some concepts of projective and affine geometry, which we use in Section 3 to write down the implicit equations of a conic arc in Bézier form. This form of writing the equations is determinant for deriving expressions for geometric characteristics of conics in Section 4.

2. Preliminaries and notation

We start by reviewing some concepts of projective geometry, that may be checked in more detail elsewhere (cfr. for instance Semple and Kneebone (1952)). Previous applications of projective geometry and point-line dual representations may be found in Farin (1999), Hoschek (1983).

The projective plane is the set of vector lines of \mathbb{R}^3 . Each *point* X of the projective plane is then characterised by a vector \vec{v} or its multiples $\lambda\vec{v}$, for any $\lambda \neq 0$.

A *frame* of the projective plane is a set $\{P, Q, R\}$ of non-aligned points. That is, if we choose vectors $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ along these lines as their respective *representatives*, they form a basis of \mathbb{R}^3 . Hence, we may write any point X of the projective plane as a combination

$$X = \alpha P + \beta Q + \gamma R, \quad (1)$$

which just states that there is a representative \vec{v} of X which satisfies

$$\vec{v} = \alpha \vec{v}_1 + \beta \vec{v}_2 + \gamma \vec{v}_3.$$

The expression 1 is unambiguous provided that the choice of vectors $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is fixed. However, the set of *coordinates* (α, β, γ) of X in this frame is equivalent to $(\lambda\alpha, \lambda\beta, \lambda\gamma)$ for $\lambda \neq 0$, since this linear combination produces a vector $\lambda\vec{v}$ which is also a representative of X . Hence, the coordinates of a point X in a frame are defined up to a non-zero multiplicative factor.

Once a frame has been chosen, we may distinguish two types of points of the projective plane:

- Points with coordinates (α, β, γ) such that $\alpha + \beta + \gamma \neq 0$. We may choose another set of coordinates $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}) = (\lambda\alpha, \lambda\beta, \lambda\gamma)$, with $\lambda = 1/(\alpha + \beta + \gamma)$ so that these points are viewed as *barycentric combinations* $\tilde{\alpha}P + \tilde{\beta}Q + \tilde{\gamma}R$, $\tilde{\alpha} + \tilde{\beta} + \tilde{\gamma} = 1$, of $\{P, Q, R\}$. See Figure 1. The frame in the projective plane provides a frame in the affine plane.

Hence the plane in \mathbb{R}^3 of equation $\alpha + \beta + \gamma = 1$ is an *affine plane* and its points are written as barycentric combinations of the frame $\{P, Q, R\}$.

- Points with coordinates (α, β, γ) such that $\alpha + \beta + \gamma = 0$. It is clear that these are not points of the affine plane. These points of the projective plane form the *line at infinity*, of equation $\alpha + \beta + \gamma = 0$. They are the directions of vectors of the affine plane introduced in the previous paragraph. See Figure 1.

We use capital letters for points in the projective plane, which may be either points in the affine plane or points at infinity. By \overrightarrow{XY} we denote the vector linking the points X and Y in the affine plane.

A line l in the projective plane is a vector plane of \mathbb{R}^3 . We may assign to it a linear map l so that $l(X) = 0$ is fulfilled by points X of the projective plane




Figure 1: The affine plane as subset of the projective plane

on the line l . For simplicity in the notation, we denote then by l both the line and a linear map that vanishes at points of the line l , that is, $l(X) = 0$ for $X \in l$. A line in the projective plane is formed by its points in the affine plane and a point at the line at infinity, which is the direction of the line. We use lower case letters for lines.

We consider now conic arcs in Bézier form,

$$c(t) = \frac{c_0(1-t)^2 + 2wc_1t(1-t) + c_2t^2}{(1-t)^2 + 2wt(1-t) + t^2}, \quad t \in [0, 1],$$

defined by their control polygon, $\{c_0, c_1, c_2\}$ (in what follows we denote $P = c_0$, $Q = c_1$, $R = c_2$), and their normalised weights, $\{1, w, 1\}$. The arc starts at $P = c(0)$ and ends at $R = c(1)$. Our frame is then the control polygon of the conic arc.

We denote by p the line through P and Q , r is the line through R and Q and q is the line through P and R . This choice reflects that p , q , r are respectively the polar lines of P , Q , R , as will be shown below.

Since $c'(0) = 2w\overrightarrow{PQ}$ and $c'(1) = 2w\overrightarrow{QR}$, the conic arc is tangent to p and r at P and R respectively (see Figure 2).

The linear maps p , q , r are fixed up to a multiplicative factor, which we may fix requiring that

$$p(R) = 1, \quad q(Q) = 1, \quad r(P) = 1. \quad (2)$$

This means that $\{r, q, p\}$ is the dual basis for $\{P, Q, R\}$ since

$$p(P) = 0 = p(Q), \quad q(P) = 0 = q(R), \quad r(Q) = 0 = r(R).$$




Figure 2: Conic arc in Bézier form

To avoid lengthy expressions, we shall denote the directions of the lines p, q, r as

$$\vec{p} := \overrightarrow{QP}, \quad \vec{q} := \overrightarrow{RP}, \quad \vec{r} := \overrightarrow{QR}. \quad (3)$$

A quick way to obtain the direction of a straight line $l = \pi p + \rho q + \sigma r$, with real coefficients π, ρ, σ , is looking for a point at infinity $\alpha P + \beta Q + \gamma R$, $\alpha + \beta + \gamma = 0$ along the line,

$$0 = (\pi p + \rho q + \sigma r)(\alpha P + \beta Q + \gamma R) = \pi\gamma + \rho\beta + \sigma\alpha.$$

A simple way of writing the solution to these equations is

$$\alpha = \lambda(\rho - \pi), \quad \beta = \lambda(\pi - \sigma), \quad \gamma = \lambda(\sigma - \rho),$$

where λ is a nonvanishing factor that we may take equal to one if we are just interested in the direction of the line:

Lemma 1. *The direction of a straight line $l = \pi p + \rho q + \sigma r$ is given by*

$$(\rho - \pi)P + (\pi - \sigma)Q + (\sigma - \rho)R,$$

or in terms of \vec{p} and \vec{r} ,

$$(\rho - \pi)\vec{p} + (\sigma - \rho)\vec{r}.$$

Conversely, the equation of a straight line through two points is readily written:

Lemma 2. *The straight line l through two points, $\alpha_1 P + \beta_1 Q + \gamma_1 R$, $\alpha_2 P + \beta_2 Q + \gamma_2 R$, is given by*

$$l = \pi p + \rho q + \sigma r, \quad \pi = \begin{vmatrix} \alpha_1 & \alpha_2 \\ \beta_1 & \beta_2 \end{vmatrix}, \quad \rho = \begin{vmatrix} \gamma_1 & \gamma_2 \\ \alpha_1 & \alpha_2 \end{vmatrix}, \quad \sigma = \begin{vmatrix} \beta_1 & \beta_2 \\ \gamma_1 & \gamma_2 \end{vmatrix},$$

just finding solutions of the system of equations

$$0 = (\pi p + \rho q + \sigma r) (\alpha_i P + \beta_i Q + \gamma_i R) = \alpha_i \sigma + \beta_i \rho + \gamma_i \pi, \quad i = 1, 2.$$

The line at infinity is $z = p + q + r$, since its points are combinations $\alpha P + \beta Q + \gamma R$ with $\alpha + \beta + \gamma = 0$.

For a conic section the *polar line* (cfr. for instance Semple and Kneebone (1952) or Richter-Gebert (2011)) of a point A is defined as:

- If A lies on the conic, the polar line of A is the tangent line at A . For instance, p is the polar line of P and r is the polar line of R .
- If A lies out of the conic, we trace both tangents to the conic through A . They meet the conic at two points. The line linking both tangency points is the polar line of A . For instance, q is the polar line of Q . If we cannot draw the tangents to the conic from a point A , we may obtain its polar line a by linking two points B and D with polar lines b , d meeting at A . See Figure 3.

In what follows, the polar line of a point and the point will be denoted by the same letter in lower and upper case respectively, as we have already done in the previous paragraphs.

We may obtain the polar line a of point A analitically: If C is the symmetric bilinear form associated to the conic, so that $C(X, X) = 0$ for points X on the conic, the points X of a satisfy $0 = C(A, X)$.

3. Implicit equations of conic sections

For determining a conic section we require five points. Providing the tangents p at P and r at R amounts to four points, since P and R are then double points. There are two degenerate conics that satisfy these conditions:




Figure 3: Examples of polar lines of points

the intersecting pair of lines $C_1 = p \cup r$ of quadratic form $p \cdot r$ ($pr + rp$ as symmetric bilinear form), which fulfills the tangency conditions at P and R , and the double line $C_2 = q \cup q$ of quadratic form q^2 , for which P and R are double points.

Hence, all conics satisfying these conditions are members of a pencil of conics

$$pr + \lambda q^2, \quad \left(\frac{pr + rp}{2} + \lambda q^2 \text{ as a symmetric form} \right), \quad (4)$$

in terms of both degenerate conic sections.

It is obvious that P and R belong to the conics in the pencil, that is

$$p(P)r(P) + \lambda q(P)^2 = 0 = p(R)r(R) + \lambda q(R)^2.$$

Using the symmetric form of the pencil of conics (4), we show that a point X is on the polar line of P if and only if

$$0 = \frac{r(P)p(X) + p(P)r(X)}{2} + \lambda q(P)q(X) = \frac{p(X)}{2},$$

and hence p is the polar line of P . Similarly, we get r as the polar line of R and q as the polar line of Q ,

$$\frac{r(Q)p + p(Q)r}{2} + \lambda q(Q)q = \lambda q.$$

A fifth condition fixes the pencil parameter λ and is obviously related to the weight w , which has not appeared so far. We may check it with the help

of the shoulder point S (Farin (2002)),

$$S := c(1/2) = \frac{P/2 + wQ + R/2}{1 + w},$$

which satisfies the conic equation,

$$0 = p(S)q(S) + \lambda q(S)^2 = \frac{1/4 + \lambda w^2}{(1 + w)^2} \Rightarrow \lambda = -\frac{1}{4w^2}.$$

The $w = \infty$ case corresponds to the degenerate conic $p \cup r$, whereas the $w = 0$ case is the double line $q \cup q$.

The polar line s of the shoulder point,

$$\frac{r(S)p + p(S)r}{2} + \lambda q(S)q = \frac{1}{4(1 + w)} \left(p + r - \frac{q}{w} \right),$$

is parallel to the line q ,

$$s(\vec{q}) = \frac{1}{4(1 + w)} (p(R) - r(P)) = 0,$$

since it contains the direction \vec{q} of q (see Figure 2).

Analogously, we may think of a dual representation, exchanging the roles of points and lines: A *point* l of the dual projective plane is a line and a *line* P of the dual projective plane is the pencil of lines through the point P . A line belongs to a conic C if it is a tangent to the conic.

We look for the tangential equation of our conic arc. That is, the implicit equation for the tangent lines to the conic.

Since p and r are respectively the tangents to the conic at P and R , there are already defined four conditions for the pencil of tangential conics.

We proceed again searching for two degenerate conics satisfying these conditions:

The degenerate conic $\tilde{C}_1 = P \cup R$ of quadratic form PR (or $(PR + RP)/2$ as a symmetric form) contains both p and r ,

$$p(P)p(R) = 0 = r(P)r(R),$$

and p is the polar line of P ,

$$p(P)R + p(R)p = p,$$

and analogously r is the polar line of R .

The other degenerate conic is $\tilde{C}_2 = Q \cup Q$ of quadratic form Q^2 , since it contains both p and r ,

$$p(Q)p(Q) = 0 = r(Q)r(Q),$$

twice.

The pencil of conics tangent to p at P and to r at R is formed then by linear combinations of both degenerate conics,

$$PR + \mu Q^2, \quad \left(\frac{PR + RP}{2} + \mu QQ \text{ as a symmetric form} \right).$$

Obviously the line p ,

$$\frac{p(R)P + p(P)R}{2} + \mu p(Q)Q = \frac{P}{2},$$

is the polar line of P and r is the polar line of R and q

$$\frac{q(R)P + q(P)R}{2} + \mu q(Q)Q = \mu Q$$

is the polar line of Q .

Again, we can fix again the free parameter μ making use of the shoulder point S .

We already know that the tangent line s at the shoulder point has linear map $s = p + r - q/w$. Hence, from

$$0 = s(R)s(P) + \mu s(Q)^2 = 1 + \frac{\mu}{w^2},$$

we learn that $\mu = -w^2$.

Summarising, we have obtained the implicit equation of a conic in terms of its control polygon and weights:

Proposition 1. *The implicit equation of a conic arc defined by its control polygon, $\{P, Q, R\}$, and its normalised weights, $\{1, w, 1\}$ is given by*

$$p(X)r(X) - \frac{q(X)^2}{4w^2} = 0. \quad (5)$$

Analogously, the tangential equation of the conic is

$$x(P)x(R) - w^2 x(Q)^2 = 0, \quad (6)$$

where p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2).

These expressions allow us to write in a straightforward and elegant manner the implicit equations of conics given in terms of their control polygons and weights, without making use of the coefficients of the conic equation,

$$a_{00} + a_{01}x + a_{02}y + a_{11}x^2 + a_{12}xy + a_{22}y^2 = 0,$$

The implicit equation of the conic in barycentric coordinates referred to the frame $\{P, Q, R\}$ is the direct result of applying the implicit equation to a point $X = \alpha P + \beta Q + \gamma R$,

$$0 = p(X)r(X) - \frac{q(X)^2}{4w^2} = \alpha\gamma - \frac{\beta^2}{4w^2}.$$

Or for a straight line as a combination of p , q and r , $x = \pi p + \rho q + \sigma r$, to be a tangent line to the conic,

$$0 = x(P)x(R) - w^2x(Q)^2 = \pi\sigma - w^2\rho^2.$$

Corollary 1. *The implicit equation of a conic arc defined by its control polygon, $\{P, Q, R\}$, and its normalised weights, $\{1, w, 1\}$ in barycentric coordinates referred to the vertices of the control polygon, $\alpha P + \beta Q + \gamma R$, is given by*

$$\alpha\gamma - \frac{\beta^2}{4w^2} = 0. \tag{7}$$

Analogously, the equation for tangent lines to the conic in barycentric coordinates, referred to the sides of the control polygon, $\pi p + \rho q + \sigma r$, is

$$\pi\sigma - w^2\rho^2 = 0, \tag{8}$$

where p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2).

4. Geometric characteristics of conics

4.1. Centres and diameters

A first instance of geometric characteristic of a conic which we characterise without making use of Euclidean geometry is the centre.

The *centre* C is, to this end, defined as the point of intersection of the diameters of the conic. Note that the *diameters* of the conic are the polar lines of the points on the line at infinity, since the tangents to the conic at




Figure 4: The centre of the conic as intersection of diameters

the intersections with a diameter are parallel. Hence, the polar line of the centre is the line at infinity, $z = p + q + r$ (see Figure 4).

Hence using the tangential equation (6) of the conic we get the centre as the point which has the line at infinity as polar line,

$$C = \frac{z(R)P + z(P)R}{2} - w^2 z(Q)Q = \frac{P + R}{2} - w^2 Q,$$

or writing it as a barycentric combination,

$$C = \frac{P + R - 2w^2 Q}{2 - 2w^2},$$

for $w \neq 1$. The case $w = 1$ corresponds to the parabola. Another simple reasoning for calculating the centre of a conic is found in Lee (1987).

Diameters are straight lines passing through the centre. Hence a line $\pi p + \rho q + \sigma r$ is a diameter if

$$0 = (\pi p + \rho q + \sigma r)(C) = \frac{\pi + \sigma - 2w^2 \rho}{2 - 2w^2}.$$

The case of the parabola, $w = 1$, can also be included here from this perspective, though within Euclidean geometry it is a noncentral conic section. In this case the centre

$$C = \frac{P + R}{2} - Q = \frac{\vec{p} + \vec{r}}{2},$$

is a direction, not a point, since it is written as a barycentric combination in which the sum of the coefficients is zero. Hence, the centre of the parabola is its only point on the line at infinity, which is the direction of the axis of the parabola. The parabola is tangent to the line at infinity at its centre. Since all diameters meet at the centre of the conic, the diameters of the parabola are straight lines parallel to the axis (see Figure 5).




Figure 5: The centre of a parabola lies at infinity

Proposition 2. *The centre of a nondegenerate conic arc defined by its control polygon, $\{P, Q, R\}$, and its normalised weights, $\{1, w, 1\}$ is given by*

$$C = \frac{P + R - 2w^2Q}{2 - 2w^2}, \quad (\vec{p} + \vec{r} \text{ if } w = 1), \quad (9)$$

and the diameters are straight lines of equation $\pi p(X) + \rho q(X) + \sigma r(X) = 0$ satisfying

$$\pi + \sigma - 2w^2\rho = 0, \quad (10)$$

where p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2).

4.2. Asymptotes

Another interesting geometric characteristic of a conic section are its *asymptotes*, defined as its tangent lines at infinity. They are real for hyperbolas and complex for ellipses. Parabolas just have a double asymptote which is the line at infinity. Again, since they are not an Euclidean feature

of conic sections, they have a simple derivation within projective and affine geometry. A different approach to calculate them is used in Goldman and Wang (2004). More on asymptotes of rational Bézier curves may be found in Farin (1999).

We get the directions of the asymptotes as the intersections of the conic with the line at infinity, searching for solutions

$$\vec{v} = \alpha P + \beta Q + \gamma R, \quad \alpha + \beta + \gamma = 0,$$

of equation (7). Taking $\beta = -2w$ to avoid denominators,

$$\begin{aligned} \vec{v}_{\pm} &= \left(w \pm \sqrt{w^2 - 1}\right) P - 2wQ + \left(w \mp \sqrt{w^2 - 1}\right) R \\ &= \left(w \pm \sqrt{w^2 - 1}\right) \vec{p} + \left(w \mp \sqrt{w^2 - 1}\right) \vec{r}, \end{aligned}$$

Of course, for the parabola, $w = 1$, we get the centre twice, $\vec{v} = \vec{p} + \vec{r}$.

The asymptotes are easily computed now as the tangents or polar lines to these points at infinity,

$$0 = r(\vec{v}_{\pm})p + p(\vec{v}_{\pm})r - \frac{q(\vec{v}_{\pm})q}{2w^2} = \left(w \pm \sqrt{w^2 - 1}\right) p + \left(w \mp \sqrt{w^2 - 1}\right) r + \frac{q}{w}.$$

For the parabola we get $p+q+r = 0$ twice, the line at infinity, as expected.

Proposition 3. *The asymptotes of a nondegenerate conic are defined by its control polygon, $\{P, Q, R\}$, and its normalised weights, $\{1, w, 1\}$ have implicit equations*

$$\left(w \pm \sqrt{w^2 - 1}\right) p(X) + \left(w \mp \sqrt{w^2 - 1}\right) r(X) + \frac{q(X)}{w} = 0, \quad (11)$$

where p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2).

The directions of the asymptotes are

$$\vec{v}_{\pm} = \left(w \pm \sqrt{w^2 - 1}\right) \vec{p} + \left(w \mp \sqrt{w^2 - 1}\right) \vec{r}, \quad (12)$$

where $\vec{p} = \overrightarrow{QP}$ and $\vec{r} = \overrightarrow{QR}$.

4.3. Axes

For obtaining the axes we finally need Euclidean geometry, but we may make use of our previous results. We denote by $\langle \vec{v}, \vec{w} \rangle$ the scalar product of two vectors \vec{v}, \vec{w} .

An *axis* is a diameter which is orthogonal to the tangent lines at the intersection points with the conic.

The axis of a conic is one of the diameters $d = \pi p + \rho q + \sigma r$ satisfying equation (10),

$$\pi + \sigma - 2w^2\rho = 0.$$

Following Lemma 1, its direction is

$$\vec{v} = (\rho - \pi)\vec{p} + (\sigma - \rho)\vec{r},$$

On the other hand, a diameter d is the polar line of a point D at infinity,

$$D = \frac{d(R)P + d(P)R}{2} - w^2 d(Q)Q = \frac{\pi P + \sigma R}{2} - w^2 \rho Q = \frac{\pi \vec{p} + \sigma \vec{r}}{2},$$

which is the direction of the tangents to the conic at the points where it intersects the diameter (see Figure 4).

If these points are vertices of the conic section, the direction \vec{v} of the diameter is orthogonal to the direction D of the tangents at the vertices,

$$\begin{aligned} 0 &= \langle \pi \vec{p} + \sigma \vec{r}, (\rho - \pi)\vec{p} + (\sigma - \rho)\vec{r} \rangle \\ &= \pi(\rho - \pi)\|\vec{p}\|^2 + \rho(\sigma - \pi)\langle \vec{p}, \vec{r} \rangle + \sigma(\sigma - \rho)\|\vec{r}\|^2. \end{aligned} \quad (13)$$

We may solve (10) and remove the ambiguity in the coefficients π, ρ, σ by fixing

$$\pi - \sigma = 2, \quad \pi = w^2\rho + 1, \quad \sigma = w^2\rho - 1,$$

so that equation (13) takes the form

$$w^2(w^2 - 1)\rho^2 + \frac{2w^2(\|\vec{r}\|^2 + \|\vec{p}\|^2) - \|\vec{q}\|^2}{(\|\vec{p}\|^2 - \|\vec{r}\|^2)}\rho + 1 = 0, \quad (14)$$

which provides the two axes:

Proposition 4. *The equations of the axes of a conic defined by its control polygon, $\{P, Q, R\}$, and its normalised weights, $\{1, w, 1\}$, $w \neq 1$, are $\pi p(X) + \rho q(X) + \sigma r(X) = 0$, with coefficients given by*

$$\pi = w^2\rho + 1, \quad \sigma = w^2\rho - 1$$

and ρ is a solution of (14), except for the case of an isosceles polygon, $\|\vec{p}\| = \|\vec{r}\|$, for which the coefficients satisfy either $\pi = -\sigma = 1$, $\rho = 0$ or $\pi = \sigma = w^2$, $\rho = 1$.

The directions of the axes are

$$\vec{v}_1 = (\rho - \pi)\vec{p} + (\sigma - \rho)\vec{r}, \quad \vec{v}_2 = \pi\vec{p} + \sigma\vec{r},$$

as linear combinations of the vectors introduced in (3).

The case of the parabola is simpler, since we already now that the axis has the direction given by the centre, a point at the line at infinity $\vec{p} + \vec{r}$, and is one of the diameters $d = \pi p + \rho q + \sigma r$ satisfying (10),

$$\pi + \sigma - 2\rho = 0.$$

The diameters intersect the conic at its centre and at another point (see Figure 5) and the direction of the tangent at it is $\pi\vec{p} + \sigma\vec{r}$. If this point is the *vertex* of the parabola, V , the direction $\vec{p} + \vec{r}$ of the axis is orthogonal to the direction of the tangent at the vertex,

$$0 = \langle \pi\vec{p} + \sigma\vec{r}, \vec{p} + \vec{r} \rangle = \pi\|\vec{p}\|^2 + (\pi + \sigma)\langle \vec{p}, \vec{r} \rangle + \sigma\|\vec{r}\|^2,$$

and we may read the coefficients of the equation of the axis,

$$\pi = 2\langle \vec{r}, \vec{p} + \vec{r} \rangle, \quad \rho = \|\vec{r}\|^2 - \|\vec{p}\|^2, \quad \sigma = -2\langle \vec{p}, \vec{p} + \vec{r} \rangle,$$

which can be obtained as the case $w = 1$ of (14).

The vertex is obtained intersecting the axis with the parabola:

Proposition 5. *The equation of the axis of a parabola defined by its control polygon, $\{P, Q, R\}$ is*

$$2\langle \vec{r}, \vec{p} + \vec{r} \rangle p + (\|\vec{r}\|^2 - \|\vec{p}\|^2) q - 2\langle \vec{p}, \vec{p} + \vec{r} \rangle r = 0,$$

where p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2) and $\vec{p} = \overrightarrow{QP}$, $\vec{r} = \overrightarrow{QR}$.

The vertex of the parabola is given by

$$V = \frac{\langle \vec{r}, \vec{p} + \vec{r} \rangle^2 P + 2\langle \vec{p}, \vec{p} + \vec{r} \rangle \langle \vec{r}, \vec{p} + \vec{r} \rangle Q + \langle \vec{p}, \vec{p} + \vec{r} \rangle^2 R}{\|\vec{p} + \vec{r}\|^4}.$$

4.4. Eccentricity

Let a be the length of the semi-major axis (the one containing the foci), b the length of the semi-minor axis (the one containing the foci) and c the semi-focal distance. The *eccentricity* of a nondegenerate conic is defined as the quotient

$$e = \frac{c}{a} \begin{cases} = 0 & \text{circumference} \\ = \frac{\sqrt{a^2 - b^2}}{a} < 1 & \text{ellipse} \\ = 1 & \text{parabola} \\ = \frac{\sqrt{a^2 + b^2}}{a} > 1 & \text{hyperbola} \end{cases}$$

taking into account that for the ellipse $a^2 = b^2 + c^2$ and for the hyperbola $c^2 = a^2 + b^2$.

It is possible to use our expressions for the asymptotes in order to calculate the eccentricity of a conic section, instead of using the lengths of the axes.

If we consider a hyperbola, the angle θ (see Figure 6) between any of the asymptotes and the major axis is related to its eccentricity,

$$\cos \theta = \frac{1}{e}.$$




Figure 6: Asymptotes of a hyperbola and its eccentricity

Since the axes produce lengthy expressions in Bézier form, it is more convenient for our purposes to use the angle 2θ between the asymptotes,

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = \frac{a^2 - b^2}{c^2} = \frac{2}{e^2} - 1,$$

taking into account that we already know the asymptotic directions v_{\pm} obtained in (12),

$$\langle v_+, v_- \rangle = \|\vec{p}\|^2 + \|\vec{r}\|^2 + 2(2w^2 - 1)\langle \vec{p}, \vec{r} \rangle = \|\vec{p} - \vec{r}\|^2 + 4w^2\langle \vec{p}, \vec{r} \rangle,$$

where the sign has been chosen in order to produce the correct definition of θ and not the complementary angle.

$$\|v_{\pm}\|^2 = (2w^2 - 1)(\|\vec{p}\|^2 + \|\vec{r}\|^2) \pm 2w\sqrt{w^2 - 1}(\|\vec{p}\|^2 - \|\vec{r}\|^2) + 2\langle \vec{p}, \vec{r} \rangle,$$

$$\begin{aligned} \|v_+\|^2\|v_-\|^2 &= \|\vec{p}\|^4 + \|\vec{r}\|^4 + 4\langle \vec{p}, \vec{r} \rangle^2 + 2(8w^4 - 8w^2 + 1)\|\vec{p}\|^2\|\vec{r}\|^2 \\ &\quad + 4(2w^2 - 1)\langle \vec{p}, \vec{r} \rangle(\|\vec{p}\|^2 + \|\vec{r}\|^2) \\ &= (\|\vec{q}\|^2 - 2w^2(\|\vec{p}\|^2 + \|\vec{r}\|^2))^2 + 4w^2(1 - w^2)(\|\vec{p}\|^2 - \|\vec{r}\|^2)^2. \end{aligned}$$

Naming

$$\Delta = \frac{\|v_+\|^2\|v_-\|^2}{w^2}, \quad b = -\frac{\langle v_+, v_- \rangle}{w^2},$$

following the notation in Xu et al. (2010), we get the formulas for the angle

$$\begin{aligned} \cos 2\theta &= \frac{\langle v_+, v_- \rangle}{\|v_+\|\|v_-\|} = \frac{-b}{\sqrt{\Delta}} \\ &= \frac{\|\vec{q}\|^2 + 4w^2\langle \vec{p}, \vec{r} \rangle}{\sqrt{(\|\vec{q}\|^2 - 2w^2(\|\vec{p}\|^2 + \|\vec{r}\|^2))^2 + 4w^2(1 - w^2)(\|\vec{p}\|^2 - \|\vec{r}\|^2)^2}}, \end{aligned} \quad (15)$$

and the eccentricity of the conic section,

$$e = \sqrt{\frac{2}{1 + \cos 2\theta}} = \sqrt{\frac{2\sqrt{\Delta}}{\sqrt{\Delta} - b}}. \quad (16)$$

A similar reasoning may be applied to ellipses, with complex asymptotes instead, leads to the same result:

Proposition 6. *The eccentricity of a conic arc defined by its control polygon, $\{P, Q, R\}$, and its normalised weights, $\{1, w, 1\}$ is given by (16), where $\vec{p} = \overrightarrow{QP}$, $\vec{q} = \overrightarrow{RP}$, $\vec{r} = \overrightarrow{QR}$.*

If the conic section is a hyperbola, the angle between its asymptotes is given by (15).

4.5. Foci

A central conic has four *foci* (two real and two complex ones) which are constructed as the intersections of the *isotropic lines*, which are the tangents to the conic drawn from two complex points located at the line at infinity named *absolute* or *circular points* I, J . The polar lines of the foci are called *directrices*.

The circular points are the points at the line at infinity where all circumferences meet (Semple and Kneebone (1952), Richter-Gebert (2011)). They are also the only points of the projective plane which are invariant under translations and rotations. The invariance under translations requires that the circular points cannot be points of the affine plane, but vectors. And the invariance under rotations requires that they have zero length. A consequence is the remarkable property that they have the same coordinates in every cartesian frame. In this section the bar denotes complex conjugation and i is the imaginary unit $\sqrt{-1}$.




Figure 7: Foci and directrices of a conic

As we see in Figure 7 (note that the diagram is fictitious due to the impossibility of drawing it in the complex plane), the polar lines i, j of the circular points provide the four tangency points of the isotropic lines. We may draw six lines through couples of these points. Four of them are the directrices.

The vanishing modulus conditions allow us to write the circular points as

complex barycentric combinations of $\{P, Q, R\}$,

$$I = aP + bQ + cR, \quad J = \bar{a}P + \bar{b}Q + \bar{c}R,$$

$$a = \|\vec{r}\|e^{i\varphi/2}, \quad c = -\|\vec{p}\|e^{-i\varphi/2}, \quad b = -a - c,$$

where φ is the angle between \vec{p} and \vec{r} .

Though these expressions are valid for every system of coordinates, it is useful for practical purposes (for instance, the examples at the end of the paper) to write down the expressions of a set of coefficients a, b, c in cartesian coordinates.

Noticing that a, b and c are defined up to a factor, we can also write these coefficients as

$$a = (R_x - Q_x) + i(R_y - Q_y),$$

$$b = (P_x - R_x) + i(P_y - R_y), \quad c = (Q_x - P_x) + i(Q_y - P_y),$$

where $(P_x, P_y), (Q_x, Q_y), (R_x, R_y)$ are the coordinates of P, Q, R in any given frame.

The linear maps of the polar lines of the circular points are then

$$i = ap + cr - \frac{b}{2w^2}q, \quad j = \bar{a}p + \bar{c}r - \frac{\bar{b}}{2w^2}q,$$

in terms of the lines p, q, r .

Hence the tangency points $\alpha P + \beta Q + \gamma R$ of the isotropic lines of I to the conic section satisfy both

$$a\gamma + c\alpha - \frac{b}{2w^2}\beta = 0, \quad \alpha\gamma - \frac{\beta^2}{4w^2} = 0, \quad (17)$$

whereas those of J satisfy the complex conjugate equations.

The simplest case is that of a parabola. The polar lines of the circular points are diameters of the conic section since the circular points lie at infinity and hence they meet at the centre of the conic. But the centre of a parabola is a *point* of the parabola. Hence, the four tangency points reduce to three: twice the centre and two other points. The parabola has then just one directrix (and one focus), defined by the two non-central tangency points.

The two solutions of equations (17) for $w = 1$ are the centre $C = P + R - 2Q$ and

$$T = \frac{a}{c}P - 2Q + \frac{c}{a}R.$$

The directrix f , the line through T and \bar{T} is, following Lemma 2, $f = \pi p + \rho q + \sigma r$, with

$$\pi = 2 \left(\frac{\bar{a}}{\bar{c}} - \frac{a}{c} \right) = \frac{\|\vec{r}\|}{\|\vec{p}\|} 4i \sin \varphi, \quad \rho = \frac{\bar{a}c}{a\bar{c}} - \frac{a\bar{c}}{a\bar{c}} = -2i \sin 2\varphi,$$

$$\sigma = 2 \left(\frac{c}{a} - \frac{\bar{c}}{\bar{a}} \right) = \frac{\|\vec{r}\|}{\|\vec{p}\|} 4i \sin \varphi,$$

or multiplying it by $\|\vec{p}\| \|\vec{r}\| / 4i \sin \varphi$, in order to get simpler real coefficients,

$$f = \|\vec{r}\|^2 p - \langle \vec{p}, \vec{r} \rangle q + \|\vec{p}\|^2 r.$$

The focus F of the parabola is the point which has f as polar line:

Proposition 7. *The equation of the directrix of a parabola defined by its control polygon, $\{P, Q, R\}$ is*

$$\|\vec{r}\|^2 p(X) - \langle \vec{p}, \vec{r} \rangle q(X) + \|\vec{p}\|^2 r(X) = 0,$$

where p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2) and $\vec{p} = \overrightarrow{QP}$, $\vec{r} = \overrightarrow{QR}$.

The focus of the parabola is

$$F = \frac{\|\vec{r}\|^2 P + 2\langle \vec{p}, \vec{r} \rangle Q + \|\vec{p}\|^2 R}{\|\vec{p} + \vec{r}\|^2}.$$

The case of central conics is more involved. Taking $\beta = 2w$, $\gamma = 1/\alpha$ for simplicity, equations (17) reduce to

$$w\alpha^2 - b\alpha + wa = 0, \tag{18}$$

for the intersections of the polar line of I with the conic.

The two points T_{\pm} of intersection of the polar line of I have

$$\alpha_{\pm} = \frac{b \pm \sqrt{b^2 - 4w^2ac}}{2wc}, \quad \beta_{\pm} = 2w, \quad \gamma_{\pm} = \frac{1}{\alpha_{\pm}} = \frac{b \mp \sqrt{b^2 - 4w^2ac}}{2wa} \tag{19}$$

as barycentric coordinates.

Their complex conjugates are the coordinates of the intersections \bar{T}_{\pm} of the polar line of \bar{I} with the conic. Note that this fact relies on keeping the

same choice of the branch of the root in (19). Otherwise the roles of \bar{T}_+ and \bar{T}_- would reverse.

In Albrecht (2001) the foci are constructed using the intersections of the tangents to the conic section from the circular points.

The pairs $\{T_+, \bar{T}_+\}$, $\{T_+, \bar{T}_-\}$, $\{T_-, \bar{T}_+\}$, $\{T_-, \bar{T}_-\}$ define the four directrices of the conic section. We are interested mainly in the two real directrices.

According to Lemma 2, it is clear that the two real directrices f_\pm are the ones spanned by $\{T_+, \bar{T}_+\}$ and $\{T_-, \bar{T}_-\}$, since their expressions $f_\pm = \pi_\pm p + \rho_\pm q + \sigma_\pm r$ have coefficients

$$\pi_\pm = 2w(\alpha_\pm - \bar{\alpha}_\pm), \quad \rho_\pm = \gamma_\pm \bar{\alpha}_\pm - \alpha_\pm \bar{\gamma}_\pm = \frac{\bar{\alpha}_\pm^2 - \alpha_\pm^2}{|\alpha_\pm|^2},$$

$$\sigma_\pm = 2w(\bar{\gamma}_\pm - \gamma_\pm) = 2w \frac{\alpha_\pm - \bar{\alpha}_\pm}{|\alpha_\pm|^2},$$

which may be simplified and made real dividing them by $2(\alpha_\pm - \bar{\alpha}_\pm)/|\alpha_\pm|^2$.

Hence, the linear maps of the real directrices of the conic section are

$$f_\pm = w|\alpha_\pm|^2 p - \Re(\alpha_\pm)q + wr, \quad (20)$$

where $\Re(\alpha_\pm) = (\alpha_\pm + \bar{\alpha}_\pm)/2$ is the real part of α_\pm .

The real foci F_\pm are the points which have the directrices f_\pm as polar lines:

Proposition 8. *The equations of the real directrices of a central conic defined by its control polygon, $\{P, Q, R\}$ and its normalised weights, $\{1, w, 1\}$, are*

$$w|\alpha_\pm|^2 p(X) - \Re(\alpha_\pm)q(X) + wr(X) = 0,$$

where the coefficients α_\pm are defined in (19) and p , q and r are respectively the linear maps corresponding to the polar lines of P , Q , R , as described in (2).

The respective foci of the conic are

$$F_\pm = \frac{|\alpha_\pm|^2 P + 2w\Re(\alpha_\pm)Q + R}{|\alpha_\pm|^2 + 2w\Re(\alpha_\pm) + 1}.$$

5. Examples

We end up with some examples of application of our results to determination of geometric characteristics of conics. Though our formulas do not depend on the choice of coordinates (barycentric, cartesian...) made and are valid for all of them, we express points in usual cartesian coordinates in the affine plane. That is, the coordinates of a point are of the form $(1, x, y)$ referred to an orthogonal basis.

5.1. Example of a parabola

We consider a parabola defined by its control polygon

$$P = (1, 0, 0), \quad Q = (1, 1/2, 0), \quad R = (1, 1, 1),$$

$$\overrightarrow{QP} = (0, -1/2, 0), \quad \overrightarrow{QR} = (0, 1/2, 1),$$

$$p(x, y) = y, \quad q(x, y) = 2x - 2y, \quad r(x, y) = 1 - 2x + y.$$

The geometric characteristics of this parabola are drawn in Figure 8.




Figure 8: Geometric characteristics of a parabola

The cartesian equation of the parabola is

$$0 = p(x, y)r(x, y) - \frac{q(x, y)^2}{4} = y - x^2 = 0.$$

The *centre* (direction of the axis) of the parabola is

$$C = P + R - 2Q = (0, 0, 1).$$

The vertex of the parabola is

$$V = \frac{\langle \vec{r}, \vec{p} + \vec{r} \rangle^2 P + 2\langle \vec{p}, \vec{p} + \vec{r} \rangle \langle \vec{r}, \vec{p} + \vec{r} \rangle Q + \langle \vec{p}, \vec{p} + \vec{r} \rangle^2 R}{\|\vec{p} + \vec{r}\|^4} = (1, 0, 0).$$

The equation of the axis a is

$$2\langle \vec{r}, \vec{p} + \vec{r} \rangle p + (\|\vec{r}\|^2 - \|\vec{p}\|^2) q - 2\langle \vec{p}, \vec{p} + \vec{r} \rangle r = 2x = 0.$$

The focus of the parabola is

$$F = \frac{\|\vec{r}\|^2 P + 2\langle \vec{p}, \vec{r} \rangle Q + \|\vec{p}\|^2 R}{\|\vec{p} + \vec{r}\|^2} = \left(1, 0, \frac{1}{4}\right).$$

And the equation of the directrix d is

$$\|\vec{r}\|^2 p - \langle \vec{p}, \vec{r} \rangle q + \|\vec{p}\|^2 r = y + \frac{1}{4} = 0.$$

5.2. Example of a hyperbola

We compute the geometric characteristics of a hyperbola with control polygon

$$P = (1, 4, 0), \quad Q = (1, 4, 1), \quad R = \left(1, 5, \frac{9}{4}\right),$$

and normalised weight $w = 3\sqrt{2}/4$. Hence

$$\overrightarrow{QP} = (0, 0, -1), \quad \overrightarrow{QR} = \left(0, 1, \frac{5}{4}\right), \quad \overrightarrow{RP} = \left(0, -1, -\frac{9}{4}\right),$$

$$p(x, y) = x - 4, \quad q(x, y) = 9 + y - \frac{9x}{4}, \quad r(x, y) = \frac{5x}{4} - 4 - y,$$

which are drawn in Figure 9.

The cartesian equation of the hiperbola is

$$0 = p(x, y)r(x, y) - \frac{q(x, y)^2}{4w^2} = \frac{x^2}{8} - \frac{2y^2}{9} - 2 = 0,$$

which is the one of a hyperbola centered at the origin with axes coincident with the cartesian axes and semiaxes lengths equal to 4 and 5. We get these results within our formulation:




Figure 9: Geometric characteristics of a hiperbola

The centre of the hiperbola is

$$C = \frac{P + R - 2w^2Q}{2 - 2w^2} = (1, 0, 0).$$

The asymptotes have equations (11),

$$\frac{\sqrt{2}}{24}(3x \pm 4y) = 0.$$

The angle between the asymptotes has cosine $7/25$ and the eccentricity is $5/4$, respectively computed with (15) and (16).

The equations of the axes are $\pi p(x, y) + \rho q(x, y) + \sigma r(x, y) = 0$, with coefficients $\pi = w^2\rho + 1$ $\sigma = w^2\rho - 1$, satisfying (14),

$$9\rho^2 - 80\rho + 64 = 0 \Rightarrow \rho = 8, \frac{8}{9}$$

and hence the equations of the axes read

$$2x = 0, \quad \frac{8y}{9} = 0,$$

as expected.

The foci of the hyperbola are

$$F_{\pm} = \frac{|\alpha_{\pm}|^2 P + w(\alpha_{\pm} + \bar{\alpha}_{\pm})Q + R}{|\alpha_{\pm}|^2 + w(\alpha_{\pm} + \bar{\alpha}_{\pm}) + 1} = (1, \pm 5, 0),$$

where

$$\alpha_+ = -\frac{3\sqrt{2}}{4} - i\frac{\sqrt{2}}{12}, \quad \alpha_- = -\frac{3\sqrt{2}}{4} + i\frac{3\sqrt{2}}{4},$$

are obtained through (19).

5.3. Example of an ellipse

Finally we provide an example of an ellipse with vertices

$$P = (1, 3, 0), \quad Q = (1, 3, 5\sqrt{3}), \quad R = \left(1, -\frac{3}{2}, \frac{5\sqrt{3}}{2}\right)$$

and normalised weight $w = 1/2$. Hence

$$\overrightarrow{QP} = (0, 0, -5\sqrt{3}), \quad \overrightarrow{QR} = \left(0, -\frac{9}{2}, -\frac{5\sqrt{3}}{2}\right), \quad \overrightarrow{RP} = \left(0, \frac{9}{2}, -\frac{5\sqrt{3}}{2}\right),$$

$$p(x, y) = \frac{2}{3} - \frac{2x}{9}, \quad q(x, y) = \frac{x}{9} + \frac{\sqrt{3}y}{15} - \frac{1}{3}, \quad r(x, y) = \frac{x}{9} - \frac{\sqrt{3}y}{15} + \frac{2}{3},$$

which are drawn in Figure 10.




Figure 10: Geometric characteristics of an ellipse

The cartesian equation of the ellipse is

$$0 = p(x, y)r(x, y) - \frac{q(x, y)^2}{4w^2} = \frac{1}{3} - \frac{x^2}{27} - \frac{y^2}{75} = 0,$$

and it corresponds to an ellipse with axes coincident with the cartesian axes and semiaxes lengths equal to 3 and 5 respectively.

The centre of the ellipse is located at

$$C = \frac{P + R - 2w^2Q}{2 - 2w^2} = (1, 0, 0).$$

and the eccentricity is $4/5$ according to (16).

The equations of the axes are $\pi p(x, y) + \rho q(x, y) + \sigma r(x, y) = 0$, with coefficients $\pi = w^2\rho + 1$ $\sigma = w^2\rho - 1$, satisfying (14),

$$-3\rho^2 + 8\rho + 16 = 0 \Rightarrow \rho = -\frac{4}{3}, 4$$

and hence they read

$$-\frac{4}{9}x = 0, \quad \frac{4\sqrt{3}}{15}y = 0.$$

The foci of the ellipse are

$$F_{\pm} = \frac{|\alpha_{\pm}|^2 P + w(\alpha_{\pm} + \bar{\alpha}_{\pm})Q + R}{|\alpha_{\pm}|^2 + w(\alpha_{\pm} + \bar{\alpha}_{\pm}) + 1} = (1, 0, \pm 4),$$

where

$$\alpha_+ = -\frac{1}{2} - \frac{2\sqrt{3}}{5} - \frac{3\sqrt{3}i}{10}, \quad \alpha_- = -\frac{1}{2} + \frac{2\sqrt{3}}{5} - \frac{3\sqrt{3}i}{10}$$

are obtained through (19).

6. Conclusions

In this paper we have shown that projective and affine geometry simplify the problem of calculating geometric characteristics of conic sections given in Bézier form, since many of these geometric characteristics can be defined without resorting to Euclidean geometry. Even for geometric characteristics which cannot be defined out of Euclidean geometry, the calculations are mitigated by using affine constructions. Closed formulas for implicit equations of conic arcs in terms of their control polygons and weights, both in standard and tangential form, have been provided. They have been used to calculate the axes, centres, diameters, foci, directrices and asymptotes of the arcs.

Acknowledgments.

The present work has been supported by Spanish Ministry of Science grant MTM2011-22886.

References

- Albrecht, G., 2001. Determination of geometrical invariants of rational parametrized conic sections, in: Lyche, T., Schumaker, L. (Eds.), *Mathematical methods for curves and surfaces*, Oslo 2000, Vanderbilt University Press, Nashville. pp. 15–24.
- Farin, G., 1999. *NURBS: from projective geometry to practical use*. A K Peters Ltd., Natick, MA.
- Farin, G., 2002. *Curves and surfaces for CAGD: a practical guide*. Morgan Kaufmann Publishers Inc., San Francisco, CA, USA.
- Goldman, R., Wang, W., 2004. Using invariants to extract geometric characteristics of conic sections from rational quadratic parameterizations. *International Journal of Computational Geometry and Applications* 14, 161 – 187.
- Hoschek, J., 1983. Dual Bézier curves and surfaces, in: Barnhill, R., Boehm, W. (Eds.), *Surfaces in CAGD*, North-Holland. pp. 147–156.
- Lee, E.T.Y., 1987. The rational Bézier representation for conics, in: Farin, G. (Ed.), *Geometric Modeling: Algorithms and new trends*, SIAM. pp. 3–20.
- Piegl, L., Tiller, W., 1997. *The NURBS book* (2nd ed.). Springer-Verlag New York, Inc., New York, NY, USA.
- Richter-Gebert, J., 2011. *Perspectives on projective geometry*. Springer-Verlag, Berlin.
- Sánchez-Reyes, J., 2004. Geometric recipes for constructing Bézier conics of given centre or focus. *Computer Aided Geometric Design* 21, 111 – 116.
- Sánchez-Reyes, J., 2011. Simple determination via complex arithmetic of geometric characteristics of Bézier conics. Preprint.

Semple, J., Kneebone, G., 1952. Algebraic projective geometry. Oxford University Press, London.

Xu, C., Kim, T., Farin, G., 2010. The eccentricity of conic sections formulated as rational Bézier quadratics. *Computer Aided Geometric Design* 27, 458 – 460.